



## **HAP Criteria for Inpatient Admissions Updated**

November 2020

We've updated HAP criteria for inpatient admissions to include clinical criteria for surgical cases.

For your convenience, a copy is attached. You can also find the criteria under *Quick Links* when you log in at **hap.org**.

# HAP Criteria for Inpatient Admissions



If the inpatient criteria are not met as follows for each of the medical conditions, observation is the expected level of care.

DIAGNOSIS	CRITERIA FOR INPATIENT ADMISSION
<b>ACUTE KIDNEY INJURY</b>	<ul style="list-style-type: none"> <li>— Creatinine level exceeds twice the patient’s baseline and the upper reference range following an observation period during which intravenous fluids were administered</li> <li><b>OR</b></li> <li>— Creatinine level continues to rise following an observation period during which intravenous fluids were administered</li> <li><b>OR</b></li> <li>— Acute renal failure with persistent nausea, vomiting or diarrhea following an observation period during which intravenous fluids were administered</li> </ul>
<b>ATRIAL FIBRILLATION</b>	<ul style="list-style-type: none"> <li>— Persistent atrial fibrillation confirmed on ECG that remains uncontrolled (rate exceeds 110) despite an observation period</li> <li><b>OR</b></li> <li>— The initiation of one of the following anti-arrhythmic agents: amiodarone, disopyramide, dofetilide, sotalol, dronedarone, mexiletine or quinidine</li> </ul>
<b>CELLULITIS</b>	<ul style="list-style-type: none"> <li>— A documented increase in area of involvement or lymph involvement despite an observation period during which intravenous anti-infectives were administered</li> <li><b>OR</b></li> <li>— Orbital cellulitis and treatment with intravenous anti-infective</li> <li><b>OR</b></li> <li>— Immunocompromised patient</li> <li><b>OR</b></li> <li>— Cellulitis in proximity with an indwelling medical device</li> <li><b>OR</b></li> <li>— Systemic toxicity</li> </ul>
<b>CHOLECYSTITIS</b>	<ul style="list-style-type: none"> <li>— Persistent systemic toxicity despite supportive care and definitive procedure that occurred in an observation period</li> <li><b>OR</b></li> <li>— Gangrenous gallbladder</li> <li><b>OR</b></li> <li>— Perforated gallbladder</li> </ul>
<b>CHRONIC OBSTRUCTIVE PULMONARY DISEASE</b>	<ul style="list-style-type: none"> <li>— Failed observation period</li> <li><b>OR</b></li> <li>— Concurrent pneumonia or heart failure confirmed on imaging</li> <li><b>OR</b></li> <li>— Ventilatory support (invasive or non-invasive)</li> <li><b>OR</b></li> <li>— Persistent documented dyspnea following 3 doses of a short acting beta-agonist and at least one criteria point: O2 sat lower than 90% and the patient’s baseline, PaO2 lower than 50mmHg, accessory respiratory muscle usage, paradoxical chest wall movement or acute or progressive central cyanosis</li> </ul>

<b>DEEP VEIN THROMBOSIS</b>	— Confirmed by imaging with at least one criteria point: Pregnancy, malignancy, pulmonary embolism, active bleeding, major surgery within the last 6 weeks, history of heparin-induced thrombocytopenia, iliofemoral vein thrombosis, creatinine level exceeds 2.5, platelet count less than 50,000, severe sustained hypertension (SBP exceeds 220 or DBP exceeds 110), bilateral DVTs, GI bleed within the last 6 months or morbid obesity (BMI exceeds 40)
<b>DEHYDRATION OR GASTROENTERITIS</b>	— Persistent volume depletion with active vomiting or diarrhea following an observation period <b>OR</b> — Orthostatic changes following an observation period
<b>DIABETIC FOOT ULCER</b>	— Ischemia <b>OR</b> — Gangrene <b>OR</b> — Systemic toxicity
<b>DIABETIC KETOACIDOSIS</b>	— Plasma glucose level exceeds 250 with at least one criteria point: pH lower than 7.25, serum bicarbonate level lower than 15 or hydroxybutyrate level exceeds 4
<b>DIABETES MELLITUS - HYPERGLYCEMIA</b>	— Persistent elevated blood sugar exceeding 450 following an observation period
<b>DYSPNEA</b>	— Mechanical ventilatory support (invasive or non-invasive) without suspected or actual diagnosis of asthma, COPD, heart failure or pneumonia
<b>FRACTURE</b>	— Fracture requiring an inpatient designated surgery within 24 hours of presentation
<b>HYPERCALCEMIA</b>	— Diagnosis of hypercalcemia along with a calcium level of 12.1-13.9 with persistent symptoms (abdominal pain, mental status changes, Glasgow Coma Scale (GCS) 9-14, nausea, nephrolithiasis, or vomiting) following an observation period
<b>HYPERKALEMIA</b>	— Potassium level exceeds 6.5 <b>OR</b> — Potassium level remains from 5.5 to 6.5 despite treatment in an observation period <b>OR</b> — Potassium level exceeds 5.4 with associated ECG changes including AV dissociation, loss of P wave, multifocal PVCs, ventricular fibrillation, ventricular tachycardia or widening QRS
<b>HYPERMAGNESEMIA</b>	— Diagnosis of hypermagnesemia along with a magnesium level greater than 3.0 with persistent symptoms (mental status changes, Glasgow Coma Scale (GCS) 9-14, muscle weakness, or vomiting) following an observation period
<b>HYPERNATERMIA</b>	— Diagnosis of hypernatremia along with a sodium level of 151-160 with persistent symptoms (hyperreflexia, irritability, mental status changes, Glasgow Coma Scale (GCS) 9-14, muscle weakness, or restlessness) following an observation period
<b>HYPERTENSION</b>	— Elevated BP (SBP exceeds 180 and/or DBP exceeds 120) along with at least one criteria point: Associated acute neurological symptoms, acute coronary syndrome, acute heart failure, pregnancy, aortic dissection, recent vascular surgery, papilledema or acute kidney injury
<b>HYPOCALCEMIA</b>	— Diagnosis of hypocalcemia along with a calcium level of 5.0-7.4 with persistent symptoms (carpopedal spasm, flaccid paralysis, muscle weakness, paresthesia, perioral numbness, or tetany) following an observation period during which treatment included calcium repletion
<b>HYPOKALEMIA</b>	— Diagnosis of hypokalemia along with a potassium level less than 2.5 and without electrocardiogram changes following an observation period during which treatment included potassium repletion

<b>HYPOMAGNESEMIA</b>	— Diagnosis of hypomagnesemia along with a magnesium level of 1.0-1.4 with persistent symptoms (carpopedal spasm, clonus, hyperreflexia, malaise, nausea, tetany, or weakness) following an observation period during which treatment included magnesium repletion
<b>HYPONATREMIA</b>	— Sodium level less than 130 following a period of observation <b>OR</b> — Sodium level less than 121
<b>HYPOPHOSPHATEMIA</b>	— Diagnosis of hypophosphatemia along with a phosphate level of 1.0-1.4 with persistent muscle weakness following an observation period during which treatment included phosphate repletion
<b>HYPOVOLEMIA OR HYPOTENSION</b>	— SBP lower than 90 <u>without</u> tachycardia, tachypnea, oliguria, mental status changes, acidosis or elevated serum lactate despite adequate fluid resuscitation <b>OR</b> — SBP lower than 90 <u>with</u> tachycardia, tachypnea, oliguria, mental status changes, acidosis or elevated serum lactate
<b>ILEUS</b>	— Ileus confirmed by imaging and unresolved following treatment during an observation period
<b>NEPHROLITHIASIS</b>	— Nephrostomy tube placement
<b>NSTEMI</b>	— Diagnosis of NSTEMI with positive cardiac biomarkers along with a cardiac catheterization performed within 24 hours of presentation <b>OR</b> scheduled to be performed within 24 hours of presentation
<b>OSTEOMYELITIS</b>	— Confirmed osteomyelitis on imaging with systemic toxicity
<b>RHABDOMYOLYSIS</b>	— CPK level exceeds 5000 despite treatment during an observation period <b>OR</b> — CPK level lower than 5000 and trending up despite treatment during an observation period <b>OR</b> — CPK level exceeds 1500 despite treatment during an observation period and the patient is unable to achieve adequate oral hydration
<b>SEPSIS -SIRS</b>	— Documented diagnosis of sepsis or SIRS with at least two qSOFA criteria points <u>following an observation period</u> during which treatment included intravenous anti-infectives and a volume expander: Sustained respiratory rate of at least 22 or higher, altered mentation or SBP of at least 100 or lower <b>OR</b> — Documented diagnosis of sepsis or SIRS with at least two criteria points <u>following an observation period</u> during which treatment included intravenous anti-infectives and a volume expander: Oral temperature exceeds 38 Celsius or lower than 36 Celsius, sustained heart rate exceeds 90, respiratory rate exceeds 20, WBC exceeds 12 or lower than 4, PaCO2 lower than 32 or bands exceed 10%
<b>STROKE</b>	— Diagnosis of stroke with confirmation on imaging of ischemia, thrombus or hemorrhage <b>AND</b> at least one new neurological deficit: aphasia, ataxia, blindness, diplopia, dysarthria, dysphagia, mental status changes (excludes coma, stupor or obtundation), Glasgow Coma Scale (GCS) 9-14, paralysis, paresis, partial or total gaze palsy, sensory deficit, or visual field loss

<b>SYNCOPE</b>	<ul style="list-style-type: none"> <li>— Documented episode of syncope with at least one criteria point: Acute myocardial ischemia, acute aortic dissection, decompensated heart failure, severe pulmonary hypertension, cardiac rhythm pauses of 3 seconds or more, pre-excitation syndromes such as Wolff-Parkinson White, suspected arrhythmogenic right ventricular cardiomyopathy, aortic or mitral stenosis, 2<sup>nd</sup> or 3<sup>rd</sup> degree AV block or primary arrhythmia syndrome such as long QT syndrome, Brugada syndrome, idiopathic ventricular tachycardia or short QT syndrome</li> </ul>
<b>TIA</b>	<ul style="list-style-type: none"> <li>— Patient's diagnosis changes to stroke during the observation period for the TIA work-up</li> </ul>
<b>VAGINAL BLEEDING</b>	<ul style="list-style-type: none"> <li>— Acute severe menorrhagia with a hemoglobin level lower than 6.1 with hemodynamic instability and treatment including blood product transfusion</li> <li><b>OR</b></li> <li>— Ruptured ectopic pregnancy</li> <li><b>OR</b></li> <li>— Uterine rupture</li> <li><b>OR</b></li> <li>— Genitourinary trauma</li> <li><b>OR</b></li> <li>— Uterine arteriovenous malformation</li> </ul>

\*Baseline\* – When no baseline is provided, baseline is considered within the standard reference range. \*LOS\* - Length of stay.

### HAP CLINICAL SURGICAL CRITERIA

<p><b>WHEN THE SURGICAL PROCEDURE IS NOT ON THE CMS (CENTERS FOR MEDICARE &amp; MEDICAID SERVICES) INPATIENT LIST OR IF THE PROCEDURE IS NOT ON THE INTERQUAL INPATIENT LIST OR IF IT IS CATEGORIZED AS AN ASTERISK PROCEDURE</b></p>	<ul style="list-style-type: none"> <li>— Three or more of the following criteria: <ul style="list-style-type: none"> <li>○ Cardiovascular disease - cardiomyopathy, unstable coronary syndromes (i.e., unstable or severe angina [Canadian Class III or IV])</li> <li>○ Uncompensated chronic heart failure [CHF] [NYHA class III or IV]</li> <li>○ BMI (Body Mass Index) greater than or equal to 40</li> <li>○ Diabetes mellitus uncontrolled despite optimal medical management with a documented A1C greater than 9.0%</li> <li>○ Hypertension which is poorly controlled despite optimal medical management (described as: systolic blood pressure <math>\geq 180</math> mmHg or diastolic blood pressure <math>\geq 110</math> mmHg)</li> <li>○ Thrombocytopenia or clotting factor disorders (hemophilia/uncontrolled coagulopathy with anticipated need for transfusions)</li> <li>○ Current treatment of a malignancy</li> <li>○ Prior documented complication with anesthesia or post-operative complications</li> <li>○ ESRD (end stage renal disease) requiring dialysis.</li> </ul> </li> <li>— One of the following criteria: <ul style="list-style-type: none"> <li>○ Advanced Liver Disease (MELD score <math>&gt;8</math>)</li> <li>○ Individual is awaiting a lung or heart transplant</li> <li>○ MI (myocardial infarction) within the last 3 months</li> <li>○ CVA (cerebrovascular accident) or TIA (transient ischemic attack) within the last 3 months.</li> </ul> </li> </ul> <p><i>Per UpTo Date: The MELD score, American Society of Anesthesiologists (ASA) class, and age predicted mortality in a study of 772 patients with cirrhosis who underwent major digestive, orthopedic, or cardiovascular surgery [41]. The MELD score was the best predictor of 30- and 90-day mortality. Mortality at 30 days ranged from 6 percent (MELD score, <math>&lt;8</math>) to more than 50 percent (MELD score, <math>&gt;20</math>) and correlated linearly with the MELD score.</i></p>
---	--

This information is not intended to represent the level of benefits covered by HAP. Please refer to the Member's Subscriber Contract, Certificate of Coverage and/or applicable Benefit Rider(s). For more information, contact HAP Customer Service at 800-801-1766.

This notification applies to all lines of HAP business and is effective until revised.

**Observation:** An observation level of care is defined by CMS as “a well-defined set of specific, clinically appropriate services, which include ongoing short-term treatment, assessment, and reassessment before a decision can be made regarding whether patients will require further treatment as hospital inpatients or if they are able to be discharged from the hospital.” This determination should be rendered within 48 hours; therefore, observation stays of up to 48 hours will be reimbursed without prior authorization. Exceptional circumstances may require extension of the observation period, of which these may be retrospectively reviewed for appropriate level of care.